

Open-sea cultivation of commercial kelps in the Atlantic coast of southern Europe

César Peteiro

Ph.D. Thesis



Madrid, Spain
2015



Universidad
Rey Juan Carlos

Escuela Superior de Ciencias Experimentales y Tecnología
Departamento de Biología, Geología, Física y Química Inorgánica
Área de Biodiversidad y Conservación

Open-sea cultivation of commercial kelps in the Atlantic coast of southern Europe

A dissertation submitted by

César Peteiro García

to the

Universidad Rey Juan Carlos (URJC)

in partial fulfillment of the requirements for the degree of

Doctor of Philosophy (Ph.D.)

in

Madrid, 2015

Supervised by

Dr. María del Brezo Martínez Díaz-Caneja

and

Dr. Noemí Sánchez Ruiz

CONTENTS

Summary	12
----------------	-----------

RESUMEN EN CASTELLANO	15
------------------------------	-----------

R1. ANTECEDENTES.....	15
R1.1. Introducción a las laminarias	15
R1.2. <i>Undaria pinnatifida</i>	19
R1.3. <i>Saccharina latissima</i>	24
R1.4. Desarrollo de la maricultura de <i>Undaria</i> y <i>Saccharina</i> en Asia	27
R1.5. Experimentos de cultivo con <i>Undaria</i> y <i>Saccharina</i> en Occidente.....	32
R2. OBJETIVOS Y ESTRUCTURA DE LA TESIS.....	35
R2.1. Objetivos generales y específicos	35
R2.2. Contenido.....	36
R2.3. Lista de manuscritos.....	36
R3. METODOLOGÍA GENERAL	40
R3.1. Obtención de plántulas en laboratorio.....	40
R3.2. Ensayos de cultivo en mar	44
R3.3. Muestreo y variables estudiadas	50
R4. CONCLUSIONES GENERALES	52
R5. BIBLIOGRAFÍA.....	53

CHAPTER I. Effect of water motion on the cultivation of the commercial seaweed <i>Undaria pinnatifida</i> in a coastal bay of Galicia, Northwest Spain	67
---	-----------

1.1. ABSTRACT.....	67
1.2. INTRODUCTION	68
1.3. MATERIAL AND METHODS	69
1.3.1. Laboratory production of seedlings	69
1.3.2. Culture sites and environmental conditions	70
1.3.3. Field cultivation	72
1.3.4. Sampling of cultivations	73
1.3.5. Statistical analysis	75
1.4. RESULTS	75
1.4.1. Environmental conditions of the culture sites.....	75
1.4.2. Growth rates, morphological variables, and yield	76
1.5. DISCUSSION	77
1.6. ACKNOWLEDGMENTS.....	81
1.7. REFERENCES	81

CHAPTER II. Outplanting time and methodologies related to mariculture of the edible kelp <i>Undaria pinnatifida</i> in the Atlantic coast of Spain	87
---	-----------

2.1. ABSTRACT.....	87
2.2. INTRODUCTION	88
2.3. MATERIAL AND METHODS	90
2.3.1. Farm site and environmental conditions	91
2.3.2. Indoor seedling production and field cultivation	91
2.3.3. Cultivation experiment 1	94

2.3.4. Cultivation experiment 2	96
2.3.5. Statistical analysis	97
2.4. RESULTS	98
2.4.1. Environmental conditions	98
2.4.2. Cultivation experiment 1	100
2.4.3. Cultivation experiment 2	101
2.5. DISCUSSION	102
2.6. ACKNOWLEDGMENTS.....	106
2.7. REFERENCES	106

CHAPTER III. Biomass yield and morphological features of the seaweed *Saccharina latissima* cultivated at two different sites in a coastal bay in the Atlantic coast of Spain **111**

3.1. ABSTRACT	111
3.2. INTRODUCTION	112
3.3. MATERIAL AND METHODS	113
3.3.1. Study site and environmental conditions.....	113
3.3.2. Seedling production and field offshore cultivation.....	114
3.3.3. Sampling of cultivations.....	117
3.3.4. Statistical analysis	118
3.4. RESULTS AND DISCUSSION	118
3.5. ACKNOWLEDGMENTS.....	124
3.6. REFERENCES	124

CHAPTER IV. Effect of outplanting time on the commercial cultivation of the kelp *Saccharina latissima* at the southern limit in the Atlantic coast, N.W. Spain **131**

4.1. ABSTRACT	131
4.2. INTRODUCTION	132
4.3. MATERIAL AND METHODS	133
4.3.1. Indoor seedling production	133
4.3.2. Study site and environmental conditions	133
4.3.3. Offshore cultivation method	135
4.3.4. Measurement of the growth rate, yield, and 'substantiality value'..	135
4.3.5. Statistical data analysis	137
4.4. RESULTS.....	137
4.5. DISCUSSION	138
4.6. CONCLUSIONS	141
4.7. ACKNOWLEDGMENTS.....	141
4.8. REFERENCES	141

CHAPTER V. Offshore cultivation methods affects blade features of the edible seaweed *Saccharina latissima* in a bay of Galicia, Northwest Spain **145**

5.1. ABSTRACT	145
5.2. INTRODUCTION	146
5.3. MATERIAL AND METHODS	147
5.3.1. Cultivation experiments with <i>Saccharina latissima</i>	147
5.3.2. Measurement of morphological features	148
5.4. RESULTS	149
5.5. DISCUSSION	150

5.6. CONCLUSION	151
5.7. ACKNOWLEDGMENTS.....	152
5.8. REFERENCES	152

**CHAPTER VI. Open-sea cultivation by transplanting young fronds of the kelp
Saccharina latissima 157**

6.1. ABSTRACT	157
6.2. INTRODUCTION	158
6.3. MATERIAL AND METHODS	160
6.3.1. Indoor production of seedlings.....	160
6.3.2. Pre-culture in greenhouse tanks	161
6.3.3. Open-sea cultivation by the transplanting method	164
6.3.4. Biomass yield and growth measurements	165
6.3.5. Environmental conditions of culture site	165
6.4. RESULTS	166
6.5. DISCUSSION	168
6.6. ACKNOWLEDGMENTS.....	172
6.7. REFERENCES	173

DISCUSSION AND CONCLUSIONS 181

D.1. KEY FACTORS IN KELP MARICULTURE.....	181
D.1.1. Hydrodynamic environment	181
D.1.2. Time frames for culture.....	184
D.1.3. Culturing methods.....	190
D.2. INTRODUCED VS. NATIVE SPECIES IN KELP MARICULTURE.....	192
D.2.1. Economic issues	192
D.2.2. Environmental issues.....	194
D.3. CONCLUSIONS.....	197
D.4. REFERENCES.....	198

SUMMARY

Kelps are one of the most economically and ecologically important groups of seaweeds in the world. They are used mainly as human food and as an alginate source for a wide range of industries; however, they also have many other applications. In addition, these large algae play important roles as ecosystem engineers and/or foundation species (kelp forest), providing habitat, protection, and food for numerous organisms in coastal ecosystems. Most kelps are confined to northern temperate regions with relatively cold water, usually below 20°C. The southern distribution limit of almost all European species is along the Atlantic coast of the Iberian Peninsula.

Commercial kelp species in eastern Asia were traditionally collected from wild stocks, although this practice has been replaced to a great extent by kelp mariculture, which now supplies more than 80% of global production. In contrast, kelps in Europe are still wild harvested for industrial purposes. However, natural resources are limited, and populations from the Atlantic coasts of the Iberian Peninsula have declined in recent years due to climate change. The development of kelp mariculture in European countries may lead to increased production for commercial uses and, in turn, protect the kelp forest from overharvesting. The introduced kelp *Undaria pinnatifida* (wakame) and the native kelp *Saccharina latissima* (sugar kombu) are economically valuable seaweeds that have been harvested for human consumption in northern Spain. Mariculture of these edible seaweeds has generated great interest as an emerging Spanish industry due to their high demand and economic value.

This doctoral thesis provides baseline information required for cultivation of kelp species on a commercial basis along the Atlantic coast of southern Europe. More specifically, it contributes to development and implementation of methodologies suitable for mariculture of *U. pinnatifida* and *S. latissima* along the Spanish Atlantic coast. The manuscript's contents describe commercial-scale cultivation trials of *U. pinnatifida* in Galicia and *S. latissima* in Galicia and Cantabria, mainly focusing on the yield and quality of cultured sporophytes. The natural life cycle consists of the microscopic gametophyte stage and the macroscopic sporophyte stage, and kelp cultivation consists of two phases related to this life cycle. In this study, a laboratory phase provided environmental conditions necessary for the artificial production of young sporophytes from gametophytes. Young sporophytes were produced at the Spanish Institute of Oceanography (IEO) in Santander, and a subsequent farming phase involved cultivation of these sporophytes in the sea until they reached a suitable size for commercial harvesting.

The main results in this thesis contribute significantly to the knowledge necessary to develop commercial-scale cultivation of kelp species in this area. The thesis, which consists of a collection of six chapters (published articles) and a general discussion, focuses on the following main topics: **(1)** the effects of hydrodynamic conditions on kelp culture grounds in coastal bays (rías) in order to identify optimal locations for cultivation of *U. pinnatifida* and *S. latissima* [Chapters I and III, Discussion]; **(2)** the suitability of different floating rafts equipped with culture systems built using horizontal rope (long-line) or hanging rope (garland and vertical types) in sheltered and more exposed environments [Chapters I, II, III, IV, V, and VI, Discussion]; **(3)** identification of the suitable time frame (planting and harvesting period) for the mariculture of both kelp species along the Atlantic coast of southern Europe (northern Spain), and its relationship with environmental factors (seawater temperature, dissolved inorganic nitrogen, underwater irradiance, and daylength) [Chapters II, III, and IV, Discussion]; and **(4)** different methods of open-sea cultivation tested with *S. latissima*, based on practices traditionally employed for the Asian *Saccharina japonica* (two year cultivation, forced cultivation, cultivation by transplanting) [Chapters III, IV, V, and VI, Discussion]. Finally, **(5)** this thesis also discusses the development of mariculture of the introduced kelp, *U. pinnatifida*, in relation to the native kelp, *S. latissima*, from an economic standpoint (e.g., yield values and range of high value-added applications) and from an environmental point of view, taking into account the risks and/or benefits associated with cultivation of those species [Discussion].

This research revealed that mariculture of the kelps *U. pinnatifida* and *S. latissima* along the Atlantic coast of southern Europe is technically and biologically viable, as indicated by the high yields obtained. However, it is highly recommended that the native *S. latissima* be cultivated in northern Spain, as it is the most economically and environmentally advantageous species due to its high biomass yield of about 16 kg fresh weight per linear meter of rope (equivalent to more than 40 tons fresh weight per hectare of a farm). This yield can be used in many value-added applications and services, such as human food, animal feed, fertilizers, and feedstock for bioethanol production. It also is an ideal candidate for use in integrated multitrophic aquaculture.

Kelps are economically important edible seaweeds with many potential applications. As the supply from wild harvest cannot meet increasing current and future uses, methods to successfully cultivate kelp species are needed. This doctoral thesis provides baseline information required for cultivation of kelp species on a commercial basis along the Atlantic coast of southern Europe. More specifically, it contributes to development and implementation of methodologies suitable for mariculture of the kelps *Undaria pinnatifida* and *Saccharina latissima* along the Spanish Atlantic coast.

